

## References of Reviewed Studies

- Anadón, P., Martín-Rubio, M., Robles, F., Rodríguez-Lázaro, J., Utrilla, R., Vázquez, A., 2010. Variation in Sr uptake in the shell of the freshwater gastropod *Bithynia tentaculata* from Lake Arreo (northern Spain) and culture experiments. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 288, 24–34. <https://doi.org/10.1016/j.palaeo.2010.01.029>
- Andrus, C.F.T., 2011. Shell midden sclerochronology. *Quat. Sci. Rev.* 30, 2892–2905. <https://doi.org/10.1016/j.quascirev.2011.07.016>
- Apolinarska, K., Kurzawska, A., 2020. Can stable isotopes of carbon and oxygen be used to determine the origin of freshwater shells used in Neolithic ornaments from Central Europe? *Archaeol. Anthropol. Sci.* 12, 15. <https://doi.org/10/gnj8bt>
- Arnold, T.E., Brenner, M., Curtis, J.H., Dutton, A., Baker, S.M., Escobar, J.H., Ortega, C.A., 2014. Application of stable isotopes ( $\delta^{18}\text{O}$ ) to determine growth patterns of the invasive gastropod *Pomacea maculata* in Florida lakes. *Florida Scientist* 77, 126–143.
- Bailey, T.R., Lear, C.H., 2006. Testing the effect of carbonate saturation on the Sr/Ca of biogenic aragonite: A case study from the River Ehen, Cumbria, UK. *Geochem. Geophys. Geosyst.* 7, n/a–n/a. <https://doi.org/10.1029/2005GC001084>
- Bar-Yosef Mayer, D.E., Leng, M.J., Aldridge, D.C., Arrowsmith, C., Gümüş, B.A., Sloane, H.J., 2012. Modern and Early-Middle Holocene shells of the freshwater mollusc *Unio*, from Çatalhöyük in the Konya Basin, Turkey: Preliminary palaeoclimatic implications from molluscan isotope data. *J. Archaeol. Sci.* 39, 76–83. <https://doi.org/10.1016/j.jas.2011.09.003>
- Black, B.A., Dunham, J.B., Blundon, B.W., Brim-Box, J., Tepley, A.J., 2015. Long-term growth-increment chronologies reveal diverse influences of climate forcing on freshwater and forest biota in the Pacific Northwest. *Glob. Change Biol.* 21, 594–604. <https://doi.org/10.1111/gcb.12756>
- Black, B.A., Dunham, J.B., Blundon, B.W., Raggon, M.F., Zima, D., 2010. Spatial variability in growth-increment chronologies of long-lived freshwater mussels: Implications for climate impacts and reconstructions. *Écoscience* 17, 240–250. <https://doi.org/10.2980/17-3-3353>
- Błazejowski, B., Racki, G., Gieszc, P., Małkowski, K., Kin, A., Krzywiecka, K., 2013. Comparative oxygen and carbon isotopic records of Miocene and recent lacustrine unionid bivalves from Poland. *Geol. Q.* 56, 113–122. <https://doi.org/10.7306/gq.1072>
- Buskirk, B.L., Bourgeois, J., Meyer, H.W., Nesbitt, E.A., 2016. Freshwater molluscan fauna from the Florissant Formation, Colorado: Paleohydrologic reconstruction of a latest Eocene lake. *Can. J. Earth Sci.* 53, 630–643. <https://doi.org/10.1139/cjes-2015-0168>
- Butitta, V.L., Rypel, A.L., Stanley, E.H., 2021. Environmental controls on long-term growth of freshwater mussels in an oligotrophic lake. *Freshw. Sci.* 40, 316–327. <https://doi.org/10.1086/714596>
- Butler, P.G., Schöne, B.R., 2017. New research in the methods and applications of sclerochronology. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 465, 295–299. <https://doi.org/10.1016/j.palaeo.2016.11.013>
- Çakırlar, C., Şeşen, R., 2013. Reading between the lines:  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  isotopes of *Unio elongatulus* shell increments as proxies for local palaeoenvironments in Mid-Holocene northern Syria. *Archaeol. Anthropol. Sci.* 5, 85–94. <https://doi.org/10.1007/s12520-013-0125-8>
- Carilli, J., Williams, B., Schöne, B.R., Krause, R.A., Fallon, S.J., 2015. Historical contaminant records from sclerochronological archives, in: Blais, J.M., Rosen, M.R., Smol, J.P. (Eds.), *Environmental Contaminants: Using Natural Archives to Track Sources and Long-Term Trends*

of Pollution, Developments in Paleoenvironmental Research, 18. Springer Netherlands: Imprint: Springer, Dordrecht.

- Carroll, M., Romanek, C., Paddock, L., 2006. The relationship between the hydrogen and oxygen isotopes of freshwater bivalve shells and their home streams. *Chem. Geol.* 234, 211–222. <https://doi.org/10.1016/j.chemgeo.2006.04.012>
- Carroll, M., Romanek, C.S., 2008. Shell layer variation in trace element concentration for the freshwater bivalve *Elliptio complanata*. *Geo-Mar. Lett.* 28, 369–381. <https://doi.org/10.1007/s00367-008-0117-3>
- Chen, F., Feng, J.-L., Ban, F.-M., Cai, B.-G., Hu, H.-P., Pei, L.-L., Wang, K.-Y., Gao, S.-P., Zhang, J.-F., 2021. Geochemistry of modern shells of the gastropod *Radix* in the Tibetan Plateau and its implications for palaeoenvironmental reconstruction. *Quat. Sci. Rev.* 251, 106703. <https://doi.org/10.1016/j.quascirev.2020.106703>
- Chen, F., Feng, J.-L., Hu, H.-P., 2016. Relationship between the shell geochemistry of the modern aquatic gastropod *Radix* and water chemistry of lakes of the Tibetan Plateau. *Hydrobiologia* 771, 239–254. <https://doi.org/10.1007/s10750-015-2634-1>
- Collins, J., Andrus, C.F.T., Scott, R.J., Moe-Hoffman, A., Peacock, E., 2020. Refit and oxygen isotope analysis of freshwater mussel shells from the Tillar Farms Site (3DR30), southeast Arkansas. *MidCont J Archaeol.* 45, 39–63. <https://doi.org/10.1080/01461109.2019.1677999>
- Collins, J.D., 2012. Assessing mussel shell diagenesis in the modern vadose zone at Lyon's Bluff (22OK520), northeast Mississippi. *J. Archaeol. Sci.* 39, 3694–3705. <https://doi.org/10.1016/j.jas.2012.06.012>
- Demény, A., Schöll-Barna, G., Fórizs, I., Osán, J., Sümegi, P., Bajnóczi, B., 2012. Stable isotope compositions and trace element concentrations in freshwater bivalve shells (*Unio* sp.) as indicators of environmental changes at Tiszapüspöki, eastern Hungary. *Cent. Eur. Geol.* 55, 441–460. <https://doi.org/10.1556/CEuGeol.55.2012.4.5>
- Dettman, D.L., Kohn, M.J., Quade, J., Ryerson, F.J., Ojha, T.P., Hamidullah, S., 2001. Seasonal stable isotope evidence for a strong Asian monsoon throughout the past 10.7 m.y. *Geology* 29, 31–34. [https://doi.org/10.1130/0091-7613\(2001\)029<0031:SSIEFA>2.0.CO;2](https://doi.org/10.1130/0091-7613(2001)029<0031:SSIEFA>2.0.CO;2)
- Dunca, E., Mutvei, H., 2001. Comparison of microgrowth pattern in *Margaritifera margaritifera* shells from south and north Sweden. *Am. Malacol. Bulletin* 16, 239–250.
- Dunca, E., Schöne, B.R., Mutvei, H., 2005. Freshwater bivalves tell of past climates: But how clearly do shells from polluted rivers speak? *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 228, 43–57. <https://doi.org/10.1016/j.palaeo.2005.03.050>
- Dunca, E., Söderberg, H., Norrgrann, O., 2011. Shell growth and age determination in the freshwater pearl mussel *Margaritifera margaritifera* in Sweden: Natural versus limed streams. *Ferrantia* 64, 48–58.
- Fritts, A.K., Fritts, M.W., Haag, W.R., DeBoer, J.A., Casper, A.F., 2017. Freshwater mussel shells (Unionidae) chronicle changes in a North American river over the past 1000 years. *Sci. Total Environ.* 575, 199–206. <https://doi.org/10.1016/j.scitotenv.2016.09.225>
- Füllenbach, C.S., Schöne, B.R., Branscheid, R., 2014. Microstructures in shells of the freshwater gastropod *Viviparus viviparus*: A potential sensor for temperature change? *Acta Biomater.* 10, 3911–3921. <https://doi.org/10.1016/j.actbio.2014.03.030>
- Gaillard, B., Lazareth, C.E., Lestrelin, H., Dufour, E., Santos, R.V., Freitas, C.E.C., Pouilly, M., 2019. Seasonal oxygen isotope variations in freshwater bivalve shells as recorders of Amazonian rivers hydrogeochemistry. *Isotopes Environ. Health. Stud.* 55, 511–525. <https://doi.org/10.1080/10256016.2019.1666120>

- Gajurel, A.P., France-Lanord, C., Huyghe, P., Guilmette, C., Gurung, D., 2006. C and O isotope compositions of modern fresh-water mollusc shells and river waters from the Himalaya and Ganga plain. *Chem. Geol.* 233, 156–183. <https://doi.org/10.1016/j.chemgeo.2006.03.002>
- Geeza, T.J., Gillikin, D.P., Goodwin, D.H., Evans, S.D., Watters, T., Warner, N.R., 2019. Controls on magnesium, manganese, strontium, and barium concentrations recorded in freshwater mussel shells from Ohio. *Chem. Geol.* 526, 142–152. <https://doi.org/10.1016/j.chemgeo.2018.01.001>
- Geeza, T.J., Gillikin, D.P., McDevitt, B., Van Sice, K., Warner, N.R., 2018. Accumulation of Marcellus Formation oil and gas wastewater metals in freshwater mussel shells. *Environ. Sci. Technol.* 52, 10883–10892. <https://doi.org/10.1021/acs.est.8b02727>
- Geist, J., Auerswald, K., Boom, A., 2005. Stable carbon isotopes in freshwater mussel shells: Environmental record or marker for metabolic activity? *Geochim. Cosmochim. Acta* 69, 3545–3554. <https://doi.org/10.1016/j.gca.2005.03.010>
- Gillikin, D.P., Hutchinson, K.A., Kumai, Y., 2009. Ontogenic increase of metabolic carbon in freshwater mussel shells (*Pyganodon cataracta*). *J. Geophys. Res.* 114, G01007. <https://doi.org/10.1029/2008JG000829>
- Gillikin, D.P., Wanamaker, A.D., Andrus, C.F.T., 2019. Chemical sclerochronology. *Chem. Geol.* 526, 1–6. <https://doi.org/10.1016/j.chemgeo.2019.06.016>
- Goewert, A., Surge, D., Carpenter, S.J., Downing, J., 2007. Oxygen and carbon isotope ratios of *Lampsilis cardium* (Unionidae) from two streams in agricultural watersheds of Iowa, USA. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 252, 637–648. <https://doi.org/10.1016/j.palaeo.2007.06.002>
- Goodwin, D.H., Gillikin, D.P., Banker, R., Watters, G.T., Dettman, D.L., Romanek, C.S., 2019. Reconstructing intra-annual growth of freshwater mussels using oxygen isotopes. *Chem. Geol.* 526, 7–22. <https://doi.org/10.1016/j.chemgeo.2018.07.030>
- Graniero, L.E., Gillikin, D.P., Surge, D., Kelemen, Z., Bouillon, S., 2021. Assessing  $\delta^{15}\text{N}$  values in the carbonate-bound organic matrix and periostracum of bivalve shells as environmental archives. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 564, 110108. <https://doi.org/10.1016/j.palaeo.2020.110108>
- Gröcke, D.R., Gillikin, D.P., 2008. Advances in mollusc sclerochronology and sclerochemistry: Tools for understanding climate and environment. *Geo-Mar. Lett.* 28, 265–268. <https://doi.org/10.1007/s00367-008-0108-4>
- Haag, W.R., 2009. Extreme longevity in freshwater mussels revisited: Sources of bias in age estimates derived from mark–recapture experiments. *Freshw. Biol.* 54, 1474–1486. <https://doi.org/10.1111/j.1365-2427.2009.02197.x>
- Haag, W.R., Rypel, A.L., 2011. Growth and longevity in freshwater mussels: Evolutionary and conservation implications. *Biol. Rev.* 86, 225–247. <https://doi.org/10.1111/j.1469-185X.2010.00146.x>
- Harzhauser, M., Mandic, O., Latal, C., Kern, A., 2012. Stable isotope composition of the Miocene Dinaride Lake System deduced from its endemic mollusc fauna. *Hydrobiologia* 682, 27–46. <https://doi.org/10.1007/s10750-011-0618-3>
- Helama, S., Läänelaid, A., Tietäväinen, H., Fauria, M.M., Kukkonen, I.T., Holopainen, J., Nielsen, J.K., Valovirta, I., 2010. Late Holocene climatic variability reconstructed from incremental data from pines and pearl mussels — A multi-proxy comparison of air and subsurface temperatures: Late Holocene climatic variability reconstructed from incremental data from pines and pearl mussels. *Boreas* 39, 734–748. <https://doi.org/10.1111/j.1502-3885.2010.00165.x>

- Helama, S., Nielsen, J.K., 2008. Construction of statistically reliable sclerochronology using subfossil shells of river pearl mussel. *J. Paleolimnol.* 40, 247–261. <https://doi.org/10.1007/s10933-007-9155-5>
- Helama, S., Nielsen, J.K., Macias Fauria, M., Valovirta, I., 2009a. A fistful of shells: Amplifying sclerochronological and palaeoclimate signals from molluscan death assemblages. *Geol. Mag.* 146, 917–930. <https://doi.org/10.1017/S0016756809990033>
- Helama, S., Nielsen, J.K., Valovirta, I., 2009b. Evaluating contemporaneity and post-mortem age of malacological remains using sclerochronology and dendrochronology. *Archaeometry* 51, 861–877. <https://doi.org/10.1111/j.1475-4754.2008.00435.x>
- Helama, S., Schöne, B.R., Black, B.A., Dunca, E., 2006. Constructing long-term proxy series for aquatic environments with absolute dating control using a sclerochronological approach: Introduction and advanced applications. *Mar. Freshwater Res.* 57, 591. <https://doi.org/10.1071/MF05176>
- Helama, S., Valovirta, I., 2014. An autoecological study of annual shell growth increments in *Margaritifera Margaritifera* from Lapland, subarctic Finland. *Memoir of the Fukui Prefectural Dinosaur Museum* 13, 25–35.
- Helama, S., Valovirta, I., 2008. Ontogenetic morphometrics of individual freshwater pearl mussels (*Margaritifera margaritifera* (L.)) reconstructed from geometric conchology and trigonometric sclerochronology. *Hydrobiologia* 610, 43–53. <https://doi.org/10.1007/s10750-008-9421-1>
- Herath, D., Jacob, D.E., Jones, H., Fallon, S.J., 2018. Potential of shells of three species of eastern Australian freshwater mussels (Bivalvia: Hyriidae) as environmental proxy archives. *Mar. Freshwater Res.* 70, 255. <https://doi.org/10.1071/MF17319>
- Izumida, H., Yoshimura, T., Suzuki, A., Nakashima, R., Ishimura, T., Yasuhara, M., Inamura, A., Shikazono, N., Kawahata, H., 2011. Biological and water chemistry controls on Sr/Ca, Ba/Ca, Mg/Ca and  $\delta^{18}\text{O}$  profiles in freshwater pearl mussel *Hyriopsis* sp. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 309, 298–308. <https://doi.org/10.1016/j.palaeo.2011.06.014>
- Kaandorp, R.J.G., Vonhof, H.B., Del Busto, C., Wesselingh, F.P., Ganssen, G.M., Marmól, A.E., Romero Pittman, L., van Hinte, J.E., 2003. Seasonal stable isotope variations of the modern Amazonian freshwater bivalve *Anodontites trapesialis*. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 194, 339–354. [https://doi.org/10.1016/S0031-0182\(03\)00332-8](https://doi.org/10.1016/S0031-0182(03)00332-8)
- Kaandorp, R.J.G., Wesselingh, F.P., Vonhof, H.B., 2006. Ecological implications from geochemical records of Miocene Western Amazonian bivalves. *J. S. Am. Earth Sci.* 21, 54–74. <https://doi.org/10.1016/j.jsames.2005.07.012>
- Kelemen, Z., Gillikin, D.P., Borges, A.V., Tambwe, E., Sembaito, A.T., Mambo, T., Wabakhangazi, J.N., Yambélé, A., Stroobandt, Y., Bouillon, S., 2021. Freshwater bivalve shells as hydrologic archives in the Congo Basin. *Geochim. Cosmochim. Acta* 308, 101–117. <https://doi.org/10.1016/j.gca.2021.05.023>
- Kelemen, Z., Gillikin, D.P., Bouillon, S., 2019. Relationship between river water chemistry and shell chemistry of two tropical African freshwater bivalve species. *Chem. Geol.* 526, 130–141. <https://doi.org/10.1016/j.chemgeo.2018.04.026>
- Kelemen, Z., Gillikin, D.P., Graniero, L.E., Havel, H., Darchambeau, F., Borges, A.V., Yambélé, A., Bassirou, A., Bouillon, S., 2017. Calibration of hydroclimate proxies in freshwater bivalve shells from Central and West Africa. *Geochim. Cosmochim. Acta* 208, 41–62. <https://doi.org/10.1016/j.gca.2017.03.025>
- Kieniewicz, J.M., Smith, J.R., 2007. Hydrologic and climatic implications of stable isotope and minor element analyses of authigenic calcite silts and gastropod shells from a mid-Pleistocene pluvial

- lake, Western Desert, Egypt. Quat. Res. 68, 431–444.  
<https://doi.org/10.1016/j.yqres.2007.07.010>
- Kukulich, S., Dettman, D., 2021. Reconstructing seasonal and baseline nitrogen isotope ratios in riverine particulate matter using freshwater mussel shells. Geochem. Geophys. Geosyst. 22.  
<https://doi.org/10.1029/2020GC009239>
- Langlet, D., Alleman, L.Y., Plisnier, P.-D., Hughes, H., André, L., 2007. Manganese content records seasonal upwelling in Lake Tanganyika mussels. Biogeosciences 4, 195–203.  
<https://doi.org/10.5194/bg-4-195-2007>
- Le Callonnec, L., Gaillot, S., Belavoine, J., Pons, F., 2019. Apports de la géochimie isotopique de bivalves dulcicoles à la reconstitution paléoenvironnementale en milieu urbain: Le cas de la ville de Lyon. Archeosciences 7–26. <https://doi.org/10.4000/archeosciences.6084>
- Leng, M.J., Lewis, J.P., 2016. Oxygen isotopes in molluscan shell: Applications in environmental archaeology. Environ. Archaeol. 21, 295–306.  
<https://doi.org/10.1179/1749631414Y.00000000048>
- Leppänen, J.J., Saarinen, T., Jilbert, T., Oulasvirta, P., 2021. The analysis of freshwater pearl mussel shells using  $\mu$ -XRF (micro-x-ray fluorescence) and the applicability for environmental reconstruction. SN Appl. Sci. 3, 1. <https://doi.org/10/gn7mnh>
- Lewis, J.P., Leng, M.J., Dean, J.R., Marciniak, A., Bar-Yosef Mayer, D.E., Wu, X., 2017. Early Holocene palaeoseasonality inferred from the stable isotope composition of *Unio* shells from Çatalhöyük, Turkey. Environ. Archaeol. 22, 79–95.  
<https://doi.org/10.1080/14614103.2015.1116216>
- Li, J., Rypel, A.L., Zhang, S.Y., Luo, Y.M., Hou, G., Murphy, B.R., Xie, S.G., 2017. Growth, longevity, and climate–growth relationships of *Corbicula fluminea* (Müller, 1774) in Hongze Lake, China. Freshw. Sci. 36, 595–608. <https://doi.org/10.1086/693463>
- Licht, A., van Cappelle, M., Abels, H.A., Ladant, J.-B., Trabucho-Alexandre, J., France-Lanord, C., Donnadiou, Y., Vandenberghe, J., Rigaudier, T., Lécuyer, C., Terry Jr, D., Adriaens, R., Boura, A., Guo, Z., Soe, A.N., Quade, J., Dupont-Nivet, G., Jaeger, J.-J., 2014. Asian monsoons in a Late Eocene greenhouse world. Nature 513, 501–506. <https://doi.org/10.1038/nature13704>
- Long, K., Schneider, L., Williams, I.S., Fallon, S.J., Stuart-Williams, H., Haberle, S., 2020. A first look at oxygen isotope records from modern and Holocene-aged gastropod (*Stenomelania*) shells from Lake Kutubu, Papua New Guinea. J. Quat. Sci. 35, 457–464.  
<https://doi.org/10.1002/jqs.3188>
- Lundquist, S.P., Worthington, T.A., Aldridge, D.C., 2019. Freshwater mussels as a tool for reconstructing climate history. Ecol. Indic. 101, 11–21.  
<https://doi.org/10.1016/j.ecolind.2018.12.048>
- Markich, S.J., Jeffree, R.A., Burke, P.T., 2002. Freshwater bivalve shells as archival indicators of metal pollution from a copper–uranium mine in tropical northern Australia. Environ. Sci. Technol. 36, 821–832. <https://doi.org/10.1021/es011066c>
- Maurer, A.-F., Galer, S.J.G., Knipper, C., Beierlein, L., Nunn, E.V., Peters, D., Tütken, T., Alt, K.W., Schöne, B.R., 2012. Bioavailable  $^{87}\text{Sr}/^{86}\text{Sr}$  in different environmental samples — Effects of anthropogenic contamination and implications for isoscapes in past migration studies. Science of The Total Environment 433, 216–229. <https://doi.org/10.1016/j.scitotenv.2012.06.046>
- McConnaughey, T.A., Gillikin, D.P., 2008. Carbon isotopes in mollusk shell carbonates. Geo-Mar. Lett. 28, 287–299. <https://doi.org/10.1007/s00367-008-0116-4>



- McLeester, M., Schurr, M., 2021. Paleoclimate of the Little Ice Age to the present in the Kankakee Valley of Illinois and Indiana, USA based on  $^{18}\text{O}/^{16}\text{O}$  isotope ratios of freshwater shells. *Environ. Archaeol.* 26, 555–566. <https://doi.org/10.1080/14614103.2020.1849487>
- Mutvei, H., Westermarck, T., 2001. How environmental information can be obtained from naiad shells, in: Bauer, G., Wächtler, K. (Eds.), *Ecology and Evolution of the Freshwater Mussels Unionoida*, Ecological Studies. Springer, Berlin, Heidelberg, pp. 367–379. [https://doi.org/10.1007/978-3-642-56869-5\\_21](https://doi.org/10.1007/978-3-642-56869-5_21)
- Neubauer, T.A., Harzhauser, M., Georgopoulou, E., Wrožyna, C., 2014. Population bottleneck triggering millennial-scale morphospace shifts in endemic thermal-spring Melanopsids. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 414, 116–128. <https://doi.org/10.1016/j.palaeo.2014.08.015>
- O’Neil, D.D., Gillikin, D.P., 2015. Do freshwater mussel shells record road-salt pollution? *Sci. Rep.* 4, 7168. <https://doi.org/10.1038/srep07168>
- Peharda, M., Schöne, B.R., Black, B.A., Corrège, T., 2021. Advances of sclerochronology research in the last decade. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 570, 110371. <https://doi.org/10.1016/j.palaeo.2021.110371>
- Pfister, L., Grave, C., Beisel, J.-N., McDonnell, J.J., 2019. A global assessment of freshwater mollusk shell oxygen isotope signatures and their relation to precipitation and stream water. *Sci. Rep.* 9, 4312. <https://doi.org/10.1038/s41598-019-40369-0>
- Pfister, L., Thielen, F., Deloule, E., Valle, N., Lentzen, E., Grave, C., Beisel, J.-N., McDonnell, J.J., 2018. Freshwater pearl mussels as a stream water stable isotope recorder. *Ecohydrology* 11, e2007. <https://doi.org/10.1002/eco.2007>
- Phung, A.T., Baeyens, W., Leermakers, M., Goderis, S., Vanhaecke, F., Gao, Y., 2013. Reproducibility of laser ablation–inductively coupled plasma–mass spectrometry (LA–ICP–MS) measurements in mussel shells and comparison with micro-drill sampling and solution ICP–MS. *Talanta* 115, 6–14. <https://doi.org/10.1016/j.talanta.2013.04.019>
- Prendergast, A.L., Pryor, A.J.E., Reade, H., Stevens, R.E., 2018. Seasonal records of palaeoenvironmental change and resource use from archaeological assemblages. *J. Archaeol. Sci. Rep.* 21, 1191–1197. <https://doi.org/10.1016/j.jasrep.2018.08.006>
- Prendergast, A.L., Stevens, R.E., 2014. Molluscs (isotopes): Analyses in environmental archaeology, in: Smith, C. (Ed.), *Encyclopedia of Global Archaeology*. Springer New York, New York, NY, pp. 5010–5019. [https://doi.org/10.1007/978-1-4419-0465-2\\_2162](https://doi.org/10.1007/978-1-4419-0465-2_2162)
- Prendergast, A.L., Versteegh, E.A.A., Schöne, B.R., 2017. New research on the development of high-resolution palaeoenvironmental proxies from geochemical properties of biogenic carbonates. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 484, 1–6. <https://doi.org/10.1016/j.palaeo.2017.05.032>
- Ricken, W., Steuber, T., Freitag, H., Hirschfeld, M., Niedenzu, B., 2003. Recent and historical discharge of a large European river system — Oxygen isotopic composition of river water and skeletal aragonite of Unionidae in the Rhine. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 193, 73–86. [https://doi.org/10.1016/S0031-0182\(02\)00713-7](https://doi.org/10.1016/S0031-0182(02)00713-7)
- Risk, M., Burchell, M., de Roo, K., Nairn, R., Tubrett, M., Forsterra, G., 2010. Trace elements in bivalve shells from the Río Cruces, Chile. *Aquat. Biol.* 10, 85–97. <https://doi.org/10.3354/ab00268>
- Roy, R., Wang, Y., Jiang, S., 2019. Growth pattern and oxygen isotopic systematics of modern freshwater mollusks along an elevation transect: Implications for paleoclimate reconstruction. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 532, 109243. <https://doi.org/10.1016/j.palaeo.2019.109243>

- Rypel, A.L., Haag, W.R., Findlay, R.H., 2009. Pervasive hydrologic effects on freshwater mussels and riparian trees in southeastern floodplain ecosystems. *Wetlands* 29, 497–504. <https://doi.org/10.1672/08-109.1>
- Rypel, A.L., Haag, W.R., Findlay, R.H., 2008. Validation of annual growth rings in freshwater mussel shells using cross dating. *Can. J. Fish. Aquat. Sci.* 65, 2224–2232. <https://doi.org/10.1139/F08-129>
- Schöll-Barna, G., 2011. An isotope mass balance model for the correlation of freshwater bivalve shell (*Unio pictorum*) carbonate  $\delta^{18}\text{O}$  to climatic conditions and water  $\delta^{18}\text{O}$  in Lake Balaton (Hungary). *J. Limnol.* 70, 272–282. <https://doi.org/10.3274/jl11-70-2-15>
- Schöll-Barna, G., Demény, A., Serlegi, G., Fábíán, S., Sümegi, P., Fórizs, I., Bajnóczi, B., 2012. Climatic variability in the Late Copper Age: Stable isotope fluctuation of prehistoric *Unio pictorum* (Unionidae) shells from Lake Balaton (Hungary). *J. Paleolimnol.* 47, 87–100. <https://doi.org/10.1007/s10933-011-9561-6>
- Schöne, B.R., 2008. The curse of physiology — Challenges and opportunities in the interpretation of geochemical data from mollusk shells. *Geo-Mar. Lett.* 28, 269–285. <https://doi.org/10.1007/s00367-008-0114-6>
- Schöne, B.R., Dunca, E., Fiebig, J., Pfeiffer, M., 2005a. Mutvei's solution: An ideal agent for resolving microgrowth structures of biogenic carbonates. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 228, 149–166. <https://doi.org/10.1016/j.palaeo.2005.03.054>
- Schöne, B.R., Dunca, E., Mutvei, H., Baier, S., Fiebig, J., 2005b. Scandinavian climate since the late 18th century reconstructed from shells of bivalve mollusks. *Zeitschrift der Dtsch. Gesellschaft für Geowissenschaften* 156, 501–515. <https://doi.org/10.1127/1860-1804/2005/0156-0501>
- Schöne, B.R., Dunca, E., Mutvei, H., Norlund, U., 2004. A 217-year record of summer air temperature reconstructed from freshwater pearl mussels (*M. margaritifera*, Sweden). *Quat. Sci. Rev.* 23, 1803–1816. <https://doi.org/10.1016/j.quascirev.2004.02.017>
- Schöne, B.R., Krause, R.A., 2016. Retrospective environmental biomonitoring — Mussel Watch expanded. *Glob. Planet Change* 144, 228–251. <https://doi.org/10.1016/j.gloplacha.2016.08.002>
- Schöne, B.R., Meret, A.E., Baier, S.M., Fiebig, J., Esper, J., McDonnell, J., Pfister, L., 2020. Freshwater pearl mussels from northern Sweden serve as long-term, high-resolution stream water isotope recorders. *Hydrol. Earth Syst. Sci.* 24, 673–696. <https://doi.org/10.5194/hess-24-673-2020>
- Schöne, B.R., Page, N.A., Rodland, D.L., Fiebig, J., Baier, S., Helama, S.O., Oschmann, W., 2007. ENSO-coupled precipitation records (1959–2004) based on shells of freshwater bivalve mollusks (*Margaritifera falcata*) from British Columbia. *Int. J. Earth Sci.* 96, 525–540. <https://doi.org/10.1007/s00531-006-0109-3>
- Schöne, B.R., Surge, D.M., 2012. Part N, Revised, Volume 1, Chapter 14: Bivalve sclerochronology and geochemistry. *Treatise Online*, no. 46 0. <https://doi.org/10.17161/to.v0i0.4297>
- Shanahan, T.M., Pigati, J.S., Dettman, D.L., Quade, J., 2005. Isotopic variability in the aragonite shells of freshwater gastropods living in springs with nearly constant temperature and isotopic composition. *Geochim. Cosmochim. Acta* 69, 3949–3966. <https://doi.org/10.1016/j.gca.2005.03.049>
- Shoults-Wilson, W.A., Seymour, L., Unrine, J.M., Wisniewski, J.M., Black, M.C., 2014. Improving data resolution and statistical rigor in the analysis of bivalve shells as environmental archives. *Environ. Sci.: Processes Impacts* 16, 247–255. <https://doi.org/10.1039/C3EM00423F>

- Soldati, A.L., Jacob, D.E., Bianchi, M.M., Hajduk, A., 2010. Microstructure and polymorphism of *Diplodon chilensis patagonicus* (d'Orbigny 1835) recent shells. *Gayana* 74, 57–65.  
<https://doi.org/10.4067/S0717-65382010000100009>
- Soldati, A.L., Jacob, D.E., Glatzel, P., Swarbrick, J.C., Geck, J., 2016. Element substitution by living organisms: The case of manganese in mollusc shell aragonite. *Sci. Rep.* 6, 22514.  
<https://doi.org/10.1038/srep22514>
- Soldati, A.L., Jacob, D.E., Schöne, B.R., Bianchi, M.M., Hajduk, A., 2009. Seasonal periodicity of growth and composition in valves of *Diplodon chilensis patagonicus* (d'Orbigny, 1835). *J. Molluscan Stud.* 75, 75–85. <https://doi.org/10.1093/mollus/eyn044>
- Stevens, R.E., Metcalfe, S.E., Leng, M.J., Lamb, A.L., Sloane, H.J., Naranjo, E., González, S., 2012. Reconstruction of Late Pleistocene climate in the Valsequillo Basin (Central Mexico) through isotopic analysis of terrestrial and freshwater snails. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 319–320, 16–27. <https://doi.org/10.1016/j.palaeo.2011.12.012>
- Sun, F., Wang, Y., Jablonski, N.G., Hou, S., Ji, X., Wolff, B., Tripathi, A., Cao, J., Yang, X., 2021. Paleoenvironment of the Late Miocene Shuitangba hominoids from Yunnan, Southwest China: Insights from stable isotopes. *Chem. Geol.* 569, 120123.  
<https://doi.org/10.1016/j.chemgeo.2021.120123>
- Taft, L., Mischke, S., Wiechert, U., Leipe, C., Rajabov, I., Riedel, F., 2014. Sclerochronological oxygen and carbon isotope ratios in *Radix* (Gastropoda) shells indicate changes of glacial meltwater flux and temperature since 4,200 cal yr BP at Lake Karakul, eastern Pamirs (Tajikistan). *J. Paleolimnol.* 52, 27–41. <https://doi.org/10.1007/s10933-014-9776-4>
- Taft, L., Wiechert, U., Riedel, F., Weynell, M., Zhang, H., 2012. Sub-seasonal oxygen and carbon isotope variations in shells of modern *Radix* sp. (Gastropoda) from the Tibetan Plateau: Potential of a new archive for palaeoclimatic studies. *Quat. Sci. Rev.* 34, 44–56.  
<https://doi.org/10.1016/j.quascirev.2011.12.006>
- Taft, L., Wiechert, U., Zhang, H., Lei, G., Mischke, S., Plessen, B., Weynell, M., Winkler, A., Riedel, F., 2013. Oxygen and carbon isotope patterns archived in shells of the aquatic gastropod *Radix*: Hydrologic and climatic signals across the Tibetan Plateau in sub-monthly resolution. *Quat. Int.* 290–291, 282–298. <https://doi.org/10.1016/j.quaint.2012.10.031>
- Tazaki, K., Morii, I., 2008. Microbial immobilization of Si, Mn, Fe, and Sr ions in the nacreous layer of *Sinohyaliopsis schlegeli* and environmental factors. *Earth Sci. Front.* 15, 54–65.  
[https://doi.org/10.1016/S1872-5791\(09\)60008-7](https://doi.org/10.1016/S1872-5791(09)60008-7)
- Thomas, K.D., 2015a. Molluscs emergent, Part I: Themes and trends in the scientific investigation of mollusc shells as resources for archaeological research. *J. Archaeol. Sci.* 56, 133–140.  
<https://doi.org/10.1016/j.jas.2015.01.024>
- Thomas, K.D., 2015b. Molluscs emergent, Part II: Themes and trends in the scientific investigation of molluscs and their shells as past human resources. *J. Archaeol. Sci.* 56, 159–167.  
<https://doi.org/10.1016/j.jas.2015.01.015>
- Trofimova, T., Alexandroff, S.J., Mette, M.J., Tray, E., Butler, P.G., Campana, S.E., Harper, E.M., Johnson, A.L.A., Morrongiello, J.R., Peharda, M., Schöne, B.R., Andersson, C., Andrus, C.F.T., Black, B.A., Burchell, M., Carroll, M.L., DeLong, K.L., Gillanders, B.M., Grønkjær, P., Killam, D., Prendergast, A.L., Reynolds, D.J., Scourse, J.D., Shirai, K., Thébault, J., Trueman, C., de Winter, N., 2020. Fundamental questions and applications of sclerochronology: Community-defined research priorities. *Estuar. Coast. Shelf Sci.* 245, 106977.  
<https://doi.org/10.1016/j.ecss.2020.106977>



- Valdovinos, C., Pedreros, P., 2007. Geographic variations in shell growth rates of the mussel *Diplodon chilensis* from temperate lakes of Chile: Implications for biodiversity conservation. *Limnologia* 37, 63–75. <https://doi.org/10.1016/j.limno.2006.08.007>
- van Hardenbroek, M., Chakraborty, A., Davies, K.L., Harding, P., Heiri, O., Henderson, A.C.G., Holmes, J.A., Lasher, G.E., Leng, M.J., Panizzo, V.N., Roberts, L., Schilder, J., Trueman, C.N., Wooller, M.J., 2018. The stable isotope composition of organic and inorganic fossils in lake sediment records: Current understanding, challenges, and future directions. *Quat. Sci. Rev.* 196, 154–176. <https://doi.org/10.1016/j.quascirev.2018.08.003>
- Van Plantinga, A.A., Grossman, E.L., 2018. Stable and clumped isotope sclerochronologies of mussels from the Brazos River, Texas (USA): Environmental and ecologic proxy. *Chem. Geol.* 502, 55–65. <https://doi.org/10.1016/j.chemgeo.2018.10.012>
- Versteegh, E.A.A., Troelstra, S.R., Vonhof, H.B., Kroon, D., 2009. Oxygen isotope composition of bivalve seasonal growth increments and ambient water in the rivers Rhine and Meuse. *PALAIOS* 24, 497–504. <https://doi.org/200908141432>
- Versteegh, E.A.A., Vonhof, H.B., Troelstra, S.R., Kaandorp, R.J.G., Kroon, D., 2010. Seasonally resolved growth of freshwater bivalves determined by oxygen and carbon isotope shell chemistry: Freshwater bivalve intra-annual growth. *Geochem. Geophys. Geosyst.* 11, n/a-n/a. <https://doi.org/10.1029/2009GC002961>
- Versteegh, E.A.A., Vonhof, H.B., Troelstra, S.R., Kroon, D., 2011. Can shells of freshwater mussels (Unionidae) be used to estimate low summer discharge of rivers and associated droughts? *Int. J. Earth Sci.* 100, 1423–1432. <https://doi.org/10.1007/s00531-010-0551-0>
- Vonhof, H.B., Joordens, J.C.A., Noback, M.L., van der Lubbe, J.H.J.L., Feibel, C.S., Kroon, D., 2013. Environmental and climatic control on seasonal stable isotope variation of freshwater molluscan bivalves in the Turkana Basin (Kenya). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 383–384, 16–26. <https://doi.org/10.1016/j.palaeo.2013.04.022>
- Watanabe, T., Suzuki, M., Komoto, Y., Shirai, K., Yamazaki, A., 2021. Daily and annual shell growth in a long-lived freshwater bivalve as a proxy for winter snowpack. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 569, 110346. <https://doi.org/10.1016/j.palaeo.2021.110346>
- Wilson, W.A., Fritts, A.K., Fritts, M.W., Unrine, J.M., Casper, A.F., 2018a. Freshwater mussel (Unionidae) shells document the decline of trace element pollution in the regional watersheds of Chicago (Illinois, USA). *Hydrobiologia* 816, 179–196. <https://doi.org/10.1007/s10750-018-3582-3>
- Wilson, W.A., Fritts, A.K., Fritts, M.W., Unrine, J.M., Tweedy, B.N., Casper, A.F., 2018b. Freshwater mussel shells (Unionidae) describe anthropogenic changes to trace element cycling within a North American river. *Sci. Total Environ.* 616–617, 1066–1076. <https://doi.org/10.1016/j.scitotenv.2017.10.212>
- Yan, H., Chen, J., Xiao, J., 2014. A review on bivalve shell, a tool for reconstruction of paleo-climate and paleo-environment. *Chin. J. Geochem.* 33, 310–315. <https://doi.org/10.1007/s11631-014-0692-0>
- Yan, H., Lee, X., Zhou, H., Cheng, H., Peng, Y., Zhou, Z., 2009. Stable isotope composition of the modern freshwater bivalve *Corbicula fluminea*. *Geochem. J.* 43, 379–387. <https://doi.org/10.2343/geochemj.1.0035>
- Yan, H., Li, Z., Lee, X., Zhou, H., Cheng, H., Chen, J., 2012. Metabolic effects on stable carbon isotopic composition of freshwater bivalve shell *Corbicula fluminea*. *Chin. J. Geochem.* 31, 103–108. <https://doi.org/10.1007/s11631-012-0555-5>

- Yan, H., Wu, G., Li, Z., Dong, D., Zheng, J., 2010. Ontogenic decrease of  $\delta^{13}\text{C}$  in bivalve shell: Evidence of vital effect. *J. Earth Sci.* 21, 271–273. <https://doi.org/10.1007/s12583-010-0233-7>
- Yoshimura, T., Izumida, H., Nakashima, R., Ishimura, T., Shikazono, N., Kawahata, H., Suzuki, A., 2015. Stable carbon isotope values in dissolved inorganic carbon of ambient waters and shell carbonate of the freshwater pearl mussel (*Hyriopsis* sp.). *J. Paleolimnol.* 54, 37–51. <https://doi.org/10.1007/s10933-015-9834-6>
- Yoshimura, T., Nakashima, R., Suzuki, A., Tomioka, N., Kawahata, H., 2010. Oxygen and carbon isotope records of cultured freshwater pearl mussel *Hyriopsis* sp. shell from Lake Kasumigaura, Japan. *J. Paleolimnol.* 43, 437–448. <https://doi.org/10.1007/s10933-009-9341-8>
- Zaarur, S., Affek, H.P., Stein, M., 2016. Last glacial-Holocene temperatures and hydrology of the Sea of Galilee and Hula Valley from clumped isotopes in *Melanopsis* shells. *Geochim. Cosmochim. Acta* 179, 142–155. <https://doi.org/10.1016/j.gca.2015.12.034>
- Zhao, L., Schöne, B.R., Mertz-Kraus, R., 2017a. Controls on strontium and barium incorporation into freshwater bivalve shells (*Corbicula fluminea*). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 465, 386–394. <https://doi.org/10.1016/j.palaeo.2015.11.040>
- Zhao, L., Walliser, E.O., Mertz-Kraus, R., Schöne, B.R., 2017b. Unionid shells (*Hyriopsis cumingii*) record manganese cycling at the sediment-water interface in a shallow eutrophic lake in China (Lake Taihu). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 484, 97–108. <https://doi.org/10.1016/j.palaeo.2017.03.010>
- Zotin, A.A., Ieshko, E.P., 2021. Growth biorhythms of the freshwater pearl mussel *Margaritifera margaritifera* (Bivalvia, Margaritiferidae), Nemina River population (Karelia). *Biol. Bull.* 48, 306–312. <https://doi.org/10.1134/S1062359021030195>
- Zuykov, M., Pelletier, E., Harper, D.A.T., 2013. Bivalve mollusks in metal pollution studies: From bioaccumulation to biomonitoring. *Chemosphere* 93, 201–208. <https://doi.org/10.1016/j.chemosphere.2013.05.001>
- Zuykov, M., Pelletier, E., Rouleau, C., Popov, L., Fowler, S.W., Orlova, M., 2009. Autoradiographic study on the distribution of  $^{241}\text{Am}$  in the shell of the freshwater zebra mussel *Dreissena polymorpha*. *Microchim. Acta.* 167, 173–178. <https://doi.org/10.1007/s00604-009-0239-y>
- Zuykov, M., Pelletier, E., Saint-Louis, R., Checa, A., Demers, S., 2012. Biosorption of thorium on the external shell surface of bivalve mollusks: The role of shell surface microtopography. *Chemosphere* 86, 680–683. <https://doi.org/10.1016/j.chemosphere.2011.11.023>
- Zuykov, M., Schindler, M., 2019. Sclerochronology-based geochemical studies of bivalve shells: Potential vs reality. *Estonian J. Earth Sci.* 68, 37. <https://doi.org/10.3176/earth.2019.05>